AMENDMENTS TO THE CLAIMS

1. (AMENDED) A compound of Formula (I)



(I)

or a stereoisomer or a pharmaceutically acceptable salt thereof, wherein:

ring B is a cycloalkyl group ~~of 3 to 8~~ <u>having 6</u> carbon atoms wherein the cycloalkyl group is saturated ~~or partially unsaturated; or a heterocycle of 3 to 7 atoms wherein the heterocycle is saturated or partially unsaturated, the heterocycle containing a heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)$_2$-, and -N(R^4)-, the heterocycle optionally containing a -C(O)-~~; ring B being substituted with 0-2 R^5;

Z is selected from a bond, -C(O)-, -C(O)NH-, -C(S)NH-, -SO_2-, and -SO_2NH-;

~~R^{1a} and R^{1b} are independently selected from H, C_{1-4} alkyl, C_{1-4} cycloalkyl, CF_3, or alternatively, R^{1a} and R^{1b} are taken together to from =O;~~

AMENDMENTS TO THE CLAIMS

R^1 is selected from a C_{6-10} aryl group substituted with 0-5 R^6 and a 5-10 membered heteroaryl system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^6;

R^2 is selected from a C_{6-10} aryl group substituted with 0-5 R^7 and a 5-10 membered heteroaryl system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^7;

R^4 is selected from H, C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, $(CRR)_qOH$, $(CRR)_tSH$, $(CRR)_tOR^{4d}$, $(CHR)_tSR^{4d}$, $(CRR)_tNR^{4a}R^{4a}$, $(CRR)_qC(O)OH$, $(CRR)_rC(O)R^{4b}$, $(CRR)_rC(O)NR^{4a}R^{4a}$, $(CRR)_tOC(O)NR^{4a}R^{4a}$, $(CRR)_tNR^{4a}C(O)OR^{4d}$, $(CRR)_tNR^{4a}C(O)R^{4b}$, $(CRR)_rC(O)OR^{4b}$, $(CRR)_tOC(O)R^{4b}$, $(CRR)_rS(O)_pR^{4b}$, $(CRR)_rS(O)_2NR^{4a}R^{4a}$, $(CRR)_rNR^{4a}S(O)_2R^{4b}$, C_{1-6} haloalkyl, a $(CRR)_r$-C_{3-10} carbocyclic residue substituted with 0-3 R^{4e}, and a $(CHR)_r$-4-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{4e};

R^{4a}, at each occurrence, is independently selected from H, methyl substituted with 0-1 R^{4c}, C_{2-6} alkyl substituted with 0-3 R^{4e}, C_{3-8} alkenyl substituted with 0-3 R^{4e}, C_{3-8} alkynyl substituted with 0-3 R^{4e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-4 R^{4e}, and a $(CHR)_r$-4-10 membered heterocyclic system containing 1-4 heteroatoms

selected from N, O, and S, substituted with 0-2 R^{4e};

R^{4b}, at each occurrence, is selected from H, C_{1-6} alkyl substituted with 0-3 R^{4e}, C_{3-8} alkenyl substituted with 0-3 R^{4e}, C_{3-8} alkynyl substituted with 0-3 R^{4e}, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-2 R^{4e}, and a $(CHR)_r$-4-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{4e};

R^{4c} is independently selected from $-C(O)R^{4b}$, $-C(O)OR^{4d}$, $-C(O)NR^{4f}R^{4f}$, and $(CH_2)_r$phenyl;

R^{4d}, at each occurrence, is selected from methyl, CF_3, C_{1-6} alkyl substituted with 0-3 R^{4e}, C_{3-8} alkenyl substituted with 0-3 R^{4e}, C_{3-8} alkynyl substituted with 0-3 R^{4e}, and a C_{3-10} carbocyclic residue substituted with 0-3 R^{4e};

R^{4e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{4f}R^{4f}$, $-C(O)R^{4i}$, $-C(O)OR^{4j}$, $-C(O)NR^{4h}R^{4h}$, $-OC(O)NR^{4h}R^{4h}$, $-NR^{4h}C(O)NR^{4h}R^{4h}$, $-NR^{4h}C(O)OR^{4j}$, and $(CH_2)_r$phenyl;

R^{4f}, at each occurrence, is selected from H, C_{1-6} alkyl, C_{3-6} cycloalkyl, and phenyl;

R^{4h}, at each occurrence, is independently selected from H, C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, and a $(CH_2)_r$-C_{3-10} carbocyclic;

R^{4i}, at each occurrence, is selected from H, C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, and a $(CH_2)_r$-C_{3-6} carbocyclic residue;

R^{4j}, at each occurrence, is selected from CF_3, C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, and a C_{3-10} carbocyclic residue;

R^5, at each occurrence, is independently selected from H, C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CRR)_rOH$, $(CRR)_rSH$, $(CRR)_rOR^{5d}$, $(CRR)_rSR^{5d}$, $(CRR)_rNR^{5a}R^{5a}$, $(CRR)_rC(O)OH$, $(CRR)_rC(O)R^{5b}$, $(CRR)_rC(O)NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}C(O)R^{5b}$, $(CRR)_rOC(O)NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}C(O)OR^{5d}$, $(CRR)_rNR^{5a}C(O)NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}C(O)H$, $(CRR)_rC(O)OR^{5b}$, $(CRR)_rOC(O)R^{5b}$, $(CRR)_rS(O)_pR^{5b}$, $(CRR)_rS(O)_2NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}S(O)_2R^{5b}$, $(CRR)_rNR^{5a}S(O)_2$ $NR^{5a}R^{5a}$, C_{1-6} haloalkyl, a $(CRR)_r$-C_{3-10} carbocyclic residue substituted with 0-3 R^{5c}, and a $(CRR)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{5c};

R^{5a}, at each occurrence, is independently selected from H, methyl substituted with 0-1 R^{5g}, C_{2-6} alkyl

substituted with 0-2 R^{5e}, C_{3-8} alkenyl substituted with 0-2 R^{5e}, C_{3-8} alkynyl substituted with 0-2 R^{5e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-5 R^{5e}, and a $(CH_2)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{5e};

R^{5b}, at each occurrence, is selected from C_{1-6} alkyl substituted with 0-3 R^{5e}, C_{3-8} alkenyl substituted with 0-2 R^{5e}, C_{3-8} alkynyl substituted with 0-2 R^{5e}, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-2 R^{5e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{5e};

R^{5c}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, Br, I, F, $(CF_2)_rCF_3$, NO_2, CN, $(CH_2)_rNR^{5f}R^{5f}$, $(CH_2)_rOH$, $(CH_2)_rOC_{1-4}$ alkyl, $(CH_2)_rSC_{1-4}$ alkyl, $(CH_2)_rC(O)OH$, $(CH_2)_rC(O)R^{5b}$, $(CH_2)_rC(O)NR^{5f}R^{5f}$, $(CH_2)_rNR^{5f}C(O)R^{5b}$, $(CH_2)_rC(O)OC_{1-4}$ alkyl, $(CH_2)_rOC(O)R^{5b}$, $(CH_2)_rC(=NR^{5f})NR^{5f}R^{5f}$, $(CH_2)_rS(O)_pR^{5b}$, $(CH_2)_rNHC(=NR^{5f})NR^{5f}R^{5f}$, $(CH_2)_rS(O)_2NR^{5f}R^{5f}$, $(CH_2)_rNR^{5f}S(O)_2R^{5b}$, and $(CH_2)_r$phenyl substituted with 0-3 R^{5e};

R^{5d}, at each occurrence, is selected from methyl, CF_3, C_{2-6} alkyl substituted with 0-2 R^{5e}, C_{3-8} alkenyl substituted with 0-2 R^{5e}, C_{3-8} alkynyl substituted with 0-2 R^{5e}, and a C_{3-10} carbocyclic residue substituted with 0-3 R^{5e};

R^{5e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-6} cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{5f}R^{5f}$, and $(CH_2)_r$phenyl;

R^{5f}, at each occurrence, is selected from H, C_{1-6} alkyl, and C_{3-6} cycloalkyl;

R^{5g} is independently selected from $-C(O)R^{5b}$, $-C(O)OR^{5d}$, $-C(O)NR^{5f}R^{5f}$, and $(CH_2)_r$phenyl;

R, at each occurrence, is selected from H, C_{1-6} alkyl substituted with R^{5e}, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, and $(CH_2)_r$phenyl substituted with R^{5e};

R^6, at each occurrence, is selected from C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, Br, I, F, NO_2, CN, $(CR'R')_rNR^{6a}R^{6a}$, $(CR'R')_rOH$, $(CR'R')_rO(CR'R')_rR^{6d}$, $(CR'R')_rSH$, $(CR'R')_rC(O)H$, $(CR'R')_rS(CR'R')_rR^{6d}$, $(CR'R')_rSC(O)(CR'R')_rR^{6b}$, $(CR'R')_rC(O)OH$, $(CR'R')_rC(O)(CR'R')_rR^{6b}$,

$(CR'R')_rNR^{6a}R^{6a}$, $(CR'R')_rC(O)NR^{6a}R^{6a}$, $(CR'R')_rNR^{6f}C(O)(CR'R')_rR^{6b}$, $(CR'R')_rC(O)O(CR'R')_rR^{6d}$, $(CR'R')_rOC(O)(CR'R')_rR^{6b}$, $(CR'R')_rOC(O)NR^{6a}(CR'R')_rR^{6d}$, $(CR'R')_rNR^{6a}C(O)NR^{6a}(CR'R')_rR^{6d}$, $(CR'R')_rNR^{6a}C(S)NR^{6a}(CR'R')_rR^{6d}$, $(CR'R')_rNR^{6f}C(O)O(CR'R')_rR^{6b}$, $(CR'R')_rC(=NR^{6f})NR^{6a}R^{6a}$, $(CR'R')_rNHC(=NR^{6f})NR^{6f}R^{6f}$, $(CR'R')_rS(O)_p(CR'R')_rR^{6b}$, $(CR'R')_rS(O)_2NR^{6a}R^{6a}$, $(CR'R')_rNR^{6f}S(O)_2NR^{6a}R^{6a}$, $(CR'R')_rNR^{6f}S(O)_2(CR'R')_rR^{6b}$, C_{1-6} haloalkyl, C_{2-8} alkenyl substituted with 0-3 R', C_{2-8} alkynyl substituted with 0-3 R', and $(CR'R')_r$phenyl substituted with 0-3 R^{6e};

alternatively, two R^6 on adjacent atoms on R^1 may join to form a cyclic acetal;

R^{6a}, at each occurrence, is selected from H, methyl substituted with 0-1 R^{6g}, C_{2-6} alkyl substituted with 0-2 R^{6e}, C_{3-8} alkenyl substituted with 0-2 R^{6e}, C_{3-8} alkynyl substituted with 0-2 R^{6e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-5 R^{6e}, and a $(CH_2)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{6e};

R^{6b}, at each occurrence, is selected from H, C_{1-6} alkyl substituted with 0-2 R^{6e}, C_{3-8} alkenyl substituted

with 0-2 R^{6e}, C_{3-8} alkynyl substituted with 0-2 R^{6e}, a $(CH_2)_rC_{3-6}$ carbocyclic residue substituted with 0-3 R^{6e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{6e};

R^{6d}, at each occurrence, is selected from C_{3-8} alkenyl substituted with 0-2 R^{6e}, C_{3-8} alkynyl substituted with 0-2 R^{6e}, methyl, CF_3, C_{2-6} alkyl substituted with 0-3 R^{6e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-3 R^{6e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{6e};

R^{6e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{6f}R^{6f}$, and $(CH_2)_r$phenyl;

R^{6f}, at each occurrence, is selected from H, C_{1-5} alkyl, and C_{3-6} cycloalkyl, and phenyl;

R^{6g} is independently selected from $-C(O)R^{6b}$, $-C(O)OR^{6d}$, $-C(O)NR^{6f}R^{6f}$, and $(CH_2)_r$phenyl;

R^7, at each occurrence, is selected from C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl,

AMENDMENTS TO THE CLAIMS

Cl, Br, I, F, NO_2, CN, $(CR'R')_rNR^{7a}R^{7a}$, $(CR'R')_rOH$, $(CR'R')_rO(CR'R')_rR^{7d}$, $(CR'R')_rSH$, $(CR'R')_rC(O)H$, $(CR'R')_rS(CR'R')_rR^{7d}$, $(CR'R')_rC(O)OH$, $(CR'R')_rC(O)(CR'R')_rR^{7b}$, $(CR'R')_rC(O)NR^{7a}R^{7a}$, $(CR'R')_rNR^{7f}C(O)(CR'R')_rR^{7b}$, $(CR'R')_rC(O)O(CR'R')_rR^{7d}$, $(CR'R')_rOC(O)(CR'R')_rR^{7b}$, $(CR'R')_rOC(O)NR^{7a}(CR'R')_rR^{7a}$, $(CR'R')_rNR^{7a}C(O)NR^{7a}(CR'R')_rR^{7a}$, $(CR'R')_rNR^{7f}C(O)O(CR'R')_rR^{7b}$, $(CR'R')_rC(=NR^{7f})NR^{7a}R^{7a}$, $(CR'R')_rNHC(=NR^{7f})NR^{7f}R^{7f}$, $(CR'R')_rS(O)_p(CR'R')_rR^{7b}$, $(CR'R')_rS(O)_2NR^{7a}R^{7a}$, $(CR'R')_rNR^{7a}S(O)_2NR^{7a}R^{7a}$, $(CR'R')_rNR^{7f}S(O)_2(CR'R')_rR^{7b}$, C_{1-6} haloalkyl, C_{2-8} alkenyl substituted with 0-3 R', C_{2-8} alkynyl substituted with 0-3 R', and $(CR'R')_r$phenyl substituted with 0-3 R^{7e};

alternatively, two R^7 on adjacent atoms on R^2 may join to form a cyclic acetal;

R^{7a}, at each occurrence, is independently selected from H, methyl substituted with 0-1 R^{7g}, C_{2-6} alkyl substituted with 0-2 R^{7e}, C_{3-8} alkenyl substituted with 0-2 R^{7e}, C_{3-8} alkynyl substituted with 0-2 R^{7e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-5 R^{7e}, and a $(CH_2)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{7e};

R^{7b}, at each occurrence, is selected from C_{1-6} alkyl substituted with 0-2 R^{7e}, C_{3-8} alkenyl substituted with 0-2 R^{7e}, C_{3-8} alkynyl substituted with 0-2 R^{7e}, a $(CH_2)_rC_{3-6}$ carbocyclic residue substituted with 0-3 R^{7e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{7e};

R^{7d}, at each occurrence, is selected from C_{3-8} alkenyl substituted with 0-2 R^{7e}, C_{3-8} alkynyl substituted with 0-2 R^{7e}, methyl, CF_3, C_{2-6} alkyl substituted with 0-3 R^{7e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-3 R^{7e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{7e};

R^{7e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{7f}R^{7f}$, and $(CH_2)_r$phenyl;

R^{7f}, at each occurrence, is selected from H, C_{1-5} alkyl, and C_{3-6} cycloalkyl, and phenyl;

R^{7g} is independently selected from $-C(O)R^{7b}$, $-C(O)OR^{7d}$, $-C(O)NR^{7f}R^{7f}$, and $(CH_2)_r$phenyl;

AMENDMENTS TO THE CLAIMS

R′, at each occurrence, is selected from H, C_{1-6} alkyl substituted with R^{6e}, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, and $(CH_2)_r$phenyl substituted with R^{6e};

R^8 is selected from H, C_{1-4} alkyl, and C_{3-4} cycloalkyl;

R^9 is selected from, H, C_{1-4} alkyl, C_{3-4} cycloalkyl, and (CH_2)-R^1;

R^{10} and R^{10a} are independently selected from H, and C_{1-4}alkyl substituted with 0-1 R^{10b},

alternatively, R^{10} and R^{10a} can join to form a C_{3-6} cycloalkyl;

R^{10b}, at each occurrence, is independently selected from -OH, -SH, $-NR^{10c}R^{10c}$, $-C(O)NR^{10c}R^{10c}$, and $-NHC(O)R^{10c}$;

R^{10c} is selected from H, C_{1-4} alkyl and C_{3-6} cycloalkyl;

R^{11} is selected from H, C_{1-4} alkyl, $(CHR)_qOH$, $(CHR)_qSH$, $(CHR)_qOR^{11d}$, $(CHR)_qS(O)_pR^{11d}$, $(CHR)_rC(O)R^{11b}$, $(CHR)_rNR^{11a}R^{11a}$, $(CHR)_rC(O)NR^{11a}R^{11a}$, $(CHR)_rC(O)NR^{11a}OR^{11d}$, $(CHR)_qNR^{11a}C(O)R^{11b}$, $(CHR)_qNR^{11a}C(O)OR^{11d}$, $(CHR)_qOC(O)NR^{11a}R^{11a}$, $(CHR)_rC(O)OR^{11d}$, a $(CHR)_r$-C_{3-6} carbocyclic residue

substituted with 0-5 R^{11e}, and a $(CHR)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{11e};

R^{11a}, at each occurrence, is independently selected from H, C_{1-4} alkyl, C_{3-4} alkenyl, C_{3-4} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-5 R^{11e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{11e};

R^{11b}, at each occurrence, is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-2 R^{11e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{11e};

R^{11d}, at each occurrence, is independently selected from H, methyl, $-CF_3$, C_{2-4} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, a C_{3-6} carbocyclic residue substituted with 0-3 R^{11e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{11e};

R^{11e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-6} cycloalkyl, Cl, F,

Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, -O-C_{1-6} alkyl, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{11f}R^{11f}$, and $(CH_2)_r$phenyl;

R^{11f}, at each occurrence, is selected from H, C_{1-6} alkyl, and C_{3-6} cycloalkyl;

R^{12} is selected from H, C_{1-4} alkyl, $(CHR)_qOH$, $(CHR)_qSH$, $(CHR)_qOR^{12d}$, $(CHR)_qS(O)_pR^{12d}$, $(CHR)_rC(O)R^{12b}$, $(CHR)_rNR^{12a}R^{12a}$, $(CHR)_rC(O)NR^{12a}R^{12a}$, $(CHR)_rC(O)NR^{12a}OR^{12d}$, $(CHR)_qNR^{12a}C(O)R^{12b}$, $(CHR)_qNR^{12a}C(O)OR^{12d}$, $(CHR)_qOC(O)NR^{12a}R^{12a}$, $(CHR)_rC(O)OR^{12d}$, a $(CHR)_r$-C_{3-6} carbocyclic residue substituted with 0-5 R^{12e}, and a $(CHR)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{12e};

R^{12a}, at each occurrence, is independently selected from H, C_{1-4} alkyl, C_{3-4} alkenyl, C_{3-4} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-5 R^{12e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{12e};

R^{12b}, at each occurrence, is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-2 R^{12e}, and a $(CH_2)_r$-5-6 membered heterocyclic

system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{12e};

R^{12d}, at each occurrence, is independently selected from H, methyl, $-CF_3$, C_{2-4} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, a C_{3-6} carbocyclic residue substituted with 0-3 R^{12e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{12e};

R^{12e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-6} cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, $-O-C_{1-6}$ alkyl, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{12f}R^{12f}$, and $(CH_2)_r$phenyl;

R^{12f}, at each occurrence, is selected from H, C_{1-6} alkyl, and C_{3-6} cycloalkyl;

~~R^{13}, at each occurrence, is independently selected from methyl, C_{2-4} alkyl substituted with 0-1 R^{13b};~~

~~R^{13b} is selected from $-OH$, $-SH$, $-NR^{13e}R^{13e}$, $-C(O)NR^{13e}R^{13e}$, and $-NHC(O)R^{13e}$;~~

~~R^{13e} is selected from H, C_{1-4} alkyl and C_{3-6} cycloalkyl;~~

n is selected from 1 and 2;

AMENDMENTS TO THE CLAIMS

~~m is selected from 0 and 1;~~

p, at each occurrence, is independently selected from 0, 1, and 2;

q, at each occurrence, is independently selected from 1, 2, 3, and 4;

r, at each occurrence, is independently selected from 0, 1, 2, 3, and 4;

~~s, at each occurrence, is independently selected from 0 and 1;~~ and

t, at each occurrence, is independently selected from 2, 3, and 4.

2. (AMENDED) A compound claim 1, wherein

ring B is a cycloalkyl group ~~of 3 to 8~~ <u>having 6</u> carbon atoms wherein the cycloalkyl group is saturated ~~or partially unsaturated; or a heterocycle of 3 to 7 atoms wherein the heterocycle is saturated or partially unsaturated, the heterocycle containing a heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)$_2$-, and -N(R^4)-, the heterocycle optionally containing a -C(O)-~~; ring B being substituted with 0-2 R^5;

Z is selected from a bond, -C(O)-, -C(O)NH-, -C(S)NH-, -SO_2-, and -SO_2NH-;

AMENDMENTS TO THE CLAIMS

~~R^{1a} and R^{1b} are independently selected from H, C_{1-4} alkyl, C_{1-4} cycloalkyl, CF_3, or alternatively, R^{1a} and R^{1b} are taken together to from =O;~~

R^1 is selected from a C_{6-10} aryl group substituted with 0-5 R^6 and a 5-10 membered heteroaryl system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^6;

R^2 is selected from a C_{6-10} aryl group substituted with 0-5 R^7 and a 5-10 membered heteroaryl system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^7;

R^4 is selected from H, C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, $(CRR)_qOH$, $(CRR)_tSH$, $(CRR)_tOR^{4d}$, $(CHR)_tSR^{4d}$, $(CRR)_tNR^{4a}R^{4a}$, $(CRR)_qC(O)OH$, $(CRR)_rC(O)R^{4b}$, $(CRR)_rC(O)NR^{4a}R^{4a}$, $(CRR)_tOC(O)NR^{4a}R^{4a}$, $(CRR)_tNR^{4a}C(O)OR^{4d}$, $(CRR)_tNR^{4a}C(O)R^{4b}$, $(CRR)_rC(O)OR^{4b}$, $(CRR)_tOC(O)R^{4b}$, $(CRR)_rS(O)_pR^{4b}$, $(CRR)_rS(O)_2NR^{4a}R^{4a}$, $(CRR)_rNR^{4a}S(O)_2R^{4b}$, C_{1-6} haloalkyl, a $(CRR)_r$-C_{3-10} carbocyclic residue substituted with 0-3 R^{4e}, and a $(CHR)_r$-4-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{4e};

R^{4a}, at each occurrence, is independently selected from H, methyl substituted with 0-1 R^{4c}, C_{2-6} alkyl substituted with 0-3 R^{4e}, C_{3-8} alkenyl substituted

with 0-3 R^{4e}, C_{3-8} alkynyl substituted with 0-3 R^{4e}, and a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-4 R^{4e};

R^{4b}, at each occurrence, is selected from H, C_{1-6} alkyl substituted with 0-3 R^{4e}, C_{3-8} alkenyl substituted with 0-3 R^{4e}, C_{3-8} alkynyl substituted with 0-3 R^{4e}, and a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-2 R^{4e};

R^{4c} is independently selected from $-C(O)R^{4b}$, $-C(O)OR^{4d}$, $-C(O)NR^{4f}R^{4f}$, and $(CH_2)_r$phenyl;

R^{4d}, at each occurrence, is selected from methyl, CF_3, C_{1-6} alkyl substituted with 0-3 R^{4e}, C_{3-8} alkenyl substituted with 0-3 R^{4e}, C_{3-8} alkynyl substituted with 0-3 R^{4e}, and a C_{3-10} carbocyclic residue substituted with 0-3 R^{4e};

R^{4e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{4f}R^{4f}$, $-C(O)R^{4i}$, $-C(O)OR^{4j}$, $-C(O)NR^{4h}R^{4h}$, $-OC(O)NR^{4h}R^{4h}$, $-NR^{4h}C(O)NR^{4h}R^{4h}$, $-NR^{4h}C(O)OR^{4j}$, and $(CH_2)_r$phenyl;

R^{4f}, at each occurrence, is selected from H, C_{1-6} alkyl, C_{3-6} cycloalkyl, and phenyl;

R^{4h}, at each occurrence, is independently selected from H, C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, and a $(CH_2)_r$-C_{3-10} carbocyclic;

R^{4i}, at each occurrence, is selected from H, C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, and a $(CH_2)_r$-C_{3-6} carbocyclic residue;

R^{4j}, at each occurrence, is selected from CF_3, C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, and a C_{3-10} carbocyclic residue;

R^5, at each occurrence, is independently selected from H, C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CRR)_rOH$, $(CRR)_rSH$, $(CRR)_rOR^{5d}$, $(CRR)_rSR^{5d}$, $(CRR)_rNR^{5a}R^{5a}$, $(CRR)_rC(O)OH$, $(CRR)_rC(O)R^{5b}$, $(CRR)_rC(O)NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}C(O)R^{5b}$, $(CRR)_rOC(O)NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}C(O)OR^{5d}$, $(CRR)_rNR^{5a}C(O)NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}C(O)H$, $(CRR)_rC(O)OR^{5b}$, $(CRR)_rOC(O)R^{5b}$, $(CRR)_rS(O)_pR^{5b}$, $(CRR)_rS(O)_2NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}S(O)_2R^{5b}$, $(CRR)_rNR^{5a}S(O)_2$ $NR^{5a}R^{5a}$, C_{1-6} haloalkyl, a $(CRR)_r$-C_{3-10} carbocyclic residue substituted with 0-3 R^{5c}, and a $(CRR)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{5c};

R^{5a}, at each occurrence, is independently selected from H, methyl substituted with 0-1 R^{5g}, C_{2-6} alkyl substituted with 0-2 R^{5e}, C_{3-8} alkenyl substituted

with 0-2 R^{5e}, C_{3-8} alkynyl substituted with 0-2 R^{5e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-5 R^{5e}, and a $(CH_2)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{5e};

R^{5b}, at each occurrence, is selected from C_{1-6} alkyl substituted with 0-3 R^{5e}, C_{3-8} alkenyl substituted with 0-2 R^{5e}, C_{3-8} alkynyl substituted with 0-2 R^{5e}, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-2 R^{5e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{5e};

R^{5c}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, Br, I, F, $(CF_2)_rCF_3$, NO_2, CN, $(CH_2)_rNR^{5f}R^{5f}$, $(CH_2)_rOH$, $(CH_2)_rOC_{1-4}$ alkyl, $(CH_2)_rSC_{1-4}$ alkyl, $(CH_2)_rC(O)OH$, $(CH_2)_rC(O)R^{5b}$, $(CH_2)_rC(O)NR^{5f}R^{5f}$, $(CH_2)_rNR^{5f}C(O)R^{5b}$, $(CH_2)_rC(O)OC_{1-4}$ alkyl, $(CH_2)_rOC(O)R^{5b}$, $(CH_2)_rC(=NR^{5f})NR^{5f}R^{5f}$, $(CH_2)_rS(O)_pR^{5b}$, $(CH_2)_rNHC(=NR^{5f})NR^{5f}R^{5f}$, $(CH_2)_rS(O)_2NR^{5f}R^{5f}$, $(CH_2)_rNR^{5f}S(O)_2R^{5b}$, and $(CH_2)_r$phenyl substituted with 0-3 R^{5e};

R^{5d}, at each occurrence, is selected from methyl, CF_3, C_{2-6} alkyl substituted with 0-2 R^{5e}, C_{3-8} alkenyl

substituted with 0-2 R^{5e}, C_{3-8} alkynyl substituted with 0-2 R^{5e}, and a C_{3-10} carbocyclic residue substituted with 0-3 R^{5e};

R^{5e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-6} cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{5f}R^{5f}$, and $(CH_2)_r$phenyl;

R^{5f}, at each occurrence, is selected from H, C_{1-6} alkyl, and C_{3-6} cycloalkyl;

R^{5g} is independently selected from $-C(O)R^{5b}$, $-C(O)OR^{5d}$, $-C(O)NR^{5f}R^{5f}$, and $(CH_2)_r$phenyl;

R, at each occurrence, is selected from H, C_{1-6} alkyl substituted with R^{5e}, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, and $(CH_2)_r$phenyl substituted with R^{5e};

R^6, at each occurrence, is selected from C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, Br, I, F, NO_2, CN, $(CR'R')_rNR^{6a}R^{6a}$, $(CR'R')_rOH$, $(CR'R')_rO(CR'R')_rR^{6d}$, $(CR'R')_rSH$, $(CR'R')_rC(O)H$, $(CR'R')_rS(CR'R')_rR^{6d}$, $(CR'R')_rC(O)OH$, $(CR'R')_rC(O)(CR'R')_rR^{6b}$, $(CR'R')_rNR^{6a}R^{6a}$, $(CR'R')_rC(O)NR^{6a}R^{6a}$, $(CR'R')_rNR^{6f}C(O)(CR'R')_rR^{6b}$, $(CR'R')_rC(O)O(CR'R')_rR^{6d}$, $(CR'R')_rOC(O)(CR'R')_rR^{6b}$,

$(CR'R')_rOC(O)NR^{6a}(CR'R')_rR^{6d}$,
$(CR'R')_rNR^{6a}C(O)NR^{6a}(CR'R')_rR^{6d}$,
$(CR'R')_rNR^{6a}C(S)NR^{6a}(CR'R')_rR^{6d}$,
$(CR'R')_rNR^{6f}C(O)O(CR'R')_rR^{6b}$,
$(CR'R')_rC(=NR^{6f})NR^{6a}R^{6a}$, $(CR'R')_rNHC(=NR^{6f})NR^{6f}R^{6f}$,
$(CR'R')_rS(O)_p(CR'R')_rR^{6b}$, $(CR'R')_rS(O)_2NR^{6a}R^{6a}$,
$(CR'R')_rNR^{6f}S(O)_2NR^{6a}R^{6a}$,
$(CR'R')_rNR^{6f}S(O)_2(CR'R')_rR^{6b}$, C_{1-6} haloalkyl, C_{2-8} alkenyl substituted with 0-3 R', C_{2-8} alkynyl substituted with 0-3 R', and $(CR'R')_r$phenyl substituted with 0-3 R^{6e};

alternatively, two R^6 on adjacent atoms on R^1 may join to form a cyclic acetal;

R^{6a}, at each occurrence, is selected from H, methyl substituted with 0-1 R^{6g}, C_{2-6} alkyl substituted with 0-2 R^{6e}, C_{3-8} alkenyl substituted with 0-2 R^{6e}, C_{3-8} alkynyl substituted with 0-2 R^{6e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-5 R^{6e}, and a $(CH_2)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{6e};

R^{6b}, at each occurrence, is selected from H, C_{1-6} alkyl substituted with 0-2 R^{6e}, C_{3-8} alkenyl substituted with 0-2 R^{6e}, C_{3-8} alkynyl substituted with 0-2 R^{6e}, a $(CH_2)_rC_{3-6}$ carbocyclic residue substituted with 0-3 R^{6e}, and a $(CH_2)_r$-5-6 membered

heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{6e};

R^{6d}, at each occurrence, is selected from C_{3-8} alkenyl substituted with 0-2 R^{6e}, C_{3-8} alkynyl substituted with 0-2 R^{6e}, methyl, CF_3, C_{2-6} alkyl substituted with 0-3 R^{6e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-3 R^{6e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{6e};

R^{6e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{6f}R^{6f}$, and $(CH_2)_r$phenyl;

R^{6f}, at each occurrence, is selected from H, C_{1-5} alkyl, and C_{3-6} cycloalkyl, and phenyl;

R^{6g} is independently selected from $-C(O)R^{6b}$, $-C(O)OR^{6d}$, $-C(O)NR^{6f}R^{6f}$, and $(CH_2)_r$phenyl;

R^7, at each occurrence, is selected from C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, Br, I, F, NO_2, CN, $(CR'R')_rNR^{7a}R^{7a}$, $(CR'R')_rOH$, $(CR'R')_rO(CR'R')_rR^{7d}$, $(CR'R')_rSH$, $(CR'R')_rC(O)H$, $(CR'R')_rS(CR'R')_rR^{7d}$, $(CR'R')_rC(O)OH$,

$(CR'R')_rC(O)(CR'R')_rR^{7b}$, $(CR'R')_rC(O)NR^{7a}R^{7a}$, $(CR'R')_rNR^{7f}C(O)(CR'R')_rR^{7b}$, $(CR'R')_rC(O)O(CR'R')_rR^{7d}$, $(CR'R')_rOC(O)(CR'R')_rR^{7b}$, $(CR'R')_rOC(O)NR^{7a}(CR'R')_rR^{7a}$, $(CR'R')_rNR^{7a}C(O)NR^{7a}(CR'R')_rR^{7a}$, $(CR'R')_rNR^{7f}C(O)O(CR'R')_rR^{7b}$, $(CR'R')_rC(=NR^{7f})NR^{7a}R^{7a}$, $(CR'R')_rNHC(=NR^{7f})NR^{7f}R^{7f}$, $(CR'R')_rS(O)_p(CR'R')_rR^{7b}$, $(CR'R')_rS(O)_2NR^{7a}R^{7a}$, $(CR'R')_rNR^{7a}S(O)_2NR^{7a}R^{7a}$, $(CR'R')_rNR^{7f}S(O)_2(CR'R')_rR^{7b}$, C_{1-6} haloalkyl, C_{2-8} alkenyl substituted with 0-3 R', C_{2-8} alkynyl substituted with 0-3 R', and $(CR'R')_r$phenyl substituted with 0-3 R^{7e};

alternatively, two R^7 on adjacent atoms on R^2 may join to form a cyclic acetal;

R^{7a}, at each occurrence, is independently selected from H, methyl substituted with 0-1 R^{7g}, C_{2-6} alkyl substituted with 0-2 R^{7e}, C_{3-8} alkenyl substituted with 0-2 R^{7e}, C_{3-8} alkynyl substituted with 0-2 R^{7e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-5 R^{7e}, and a $(CH_2)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{7e};

R^{7b}, at each occurrence, is selected from C_{1-6} alkyl substituted with 0-2 R^{7e}, C_{3-8} alkenyl substituted

with 0-2 R^{7e}, C_{3-8} alkynyl substituted with 0-2 R^{7e}, a $(CH_2)_rC_{3-6}$ carbocyclic residue substituted with 0-3 R^{7e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-2 R^{7e};

R^{7d}, at each occurrence, is selected from C_{3-8} alkenyl substituted with 0-2 R^{7e}, C_{3-8} alkynyl substituted with 0-2 R^{7e}, methyl, CF_3, C_{2-6} alkyl substituted with 0-3 R^{7e}, a $(CH_2)_r$-C_{3-10} carbocyclic residue substituted with 0-3 R^{7e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{7e};

R^{7e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{7f}R^{7f}$, and $(CH_2)_r$phenyl;

R^{7f}, at each occurrence, is selected from H, C_{1-5} alkyl, and C_{3-6} cycloalkyl, and phenyl;

R^{7g} is independently selected from $-C(O)R^{7b}$, $-C(O)OR^{7d}$, $-C(O)NR^{7f}R^{7f}$, and $(CH_2)_r$phenyl;

R′, at each occurrence, is selected from H, C_{1-6} alkyl substituted with R^{6e}, C_{2-8} alkenyl, C_{2-8} alkynyl,

AMENDMENTS TO THE CLAIMS

$(CH_2)_rC_{3-6}$ cycloalkyl, and $(CH_2)_r$phenyl substituted with R^{6e};

R^8 is selected from H, C_{1-4} alkyl, and C_{3-4} cycloalkyl;

R^9 is selected from, H, C_{1-4} alkyl, C_{3-4} cycloalkyl, and (CH_2)-R^1;

R^{10} and R^{10a} are independently selected from H, and C_{1-4}alkyl substituted with 0-1 R^{10b},

alternatively, R^{10} and R^{10a} can join to form a C_{3-6} cycloalkyl;

R^{10b}, at each occurrence, is independently selected from -OH, -SH, $-NR^{10c}R^{10c}$, $-C(O)NR^{10c}R^{10c}$, and $-NHC(O)R^{10c}$;

R^{10c} is selected from H, C_{1-4} alkyl and C_{3-6} cycloalkyl;

R^{11} is selected from H, C_{1-4} alkyl, $(CHR)_qOH$, $(CHR)_qSH$, $(CHR)_qOR^{11d}$, $(CHR)_qS(O)_pR^{11d}$, $(CHR)_rC(O)R^{11b}$, $(CHR)_rNR^{11a}R^{11a}$, $(CHR)_rC(O)NR^{11a}R^{11a}$, $(CHR)_rC(O)NR^{11a}OR^{11d}$, $(CHR)_qNR^{11a}C(O)R^{11b}$, $(CHR)_qNR^{11a}C(O)OR^{11d}$, $(CHR)_qOC(O)NR^{11a}R^{11a}$, $(CHR)_rC(O)OR^{11d}$, a $(CHR)_r$-C_{3-6} carbocyclic residue substituted with 0-5 R^{11e}, and a $(CHR)_r$-5-10 membered heterocyclic system containing 1-4

heteroatoms selected from N, O, and S, substituted with 0-3 R^{11e};

R^{11a}, at each occurrence, is independently selected from H, C_{1-4} alkyl, C_{3-4} alkenyl, C_{3-4} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-5 R^{11e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{11e};

R^{11b}, at each occurrence, is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-2 R^{11e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{11e};

R^{11d}, at each occurrence, is independently selected from H, methyl, $-CF_3$, C_{2-4} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, a C_{3-6} carbocyclic residue substituted with 0-3 R^{11e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{11e};

R^{11e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-6} cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH,

$-O-C_{1-6}$ alkyl, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{11f}R^{11f}$, and $(CH_2)_r$phenyl;

R^{11f}, at each occurrence, is selected from H, C_{1-6} alkyl, and C_{3-6} cycloalkyl;

R^{12} is selected from H, C_{1-4} alkyl, $(CHR)_qOH$, $(CHR)_qSH$, $(CHR)_qOR^{12d}$, $(CHR)_qS(O)_pR^{12d}$, $(CHR)_rC(O)R^{12b}$, $(CHR)_rNR^{12a}R^{12a}$, $(CHR)_rC(O)NR^{12a}R^{12a}$, $(CHR)_rC(O)NR^{12a}OR^{12d}$, $(CHR)_qNR^{12a}C(O)R^{12b}$, $(CHR)_qNR^{12a}C(O)OR^{12d}$, $(CHR)_qOC(O)NR^{12a}R^{12a}$, $(CHR)_rC(O)OR^{12d}$, a $(CHR)_r$-C_{3-6} carbocyclic residue substituted with 0-5 R^{12e}, and a $(CHR)_r$-5-10 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{12e};

R^{12a}, at each occurrence, is independently selected from H, C_{1-4} alkyl, C_{3-4} alkenyl, C_{3-4} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-5 R^{12e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{12e};

R^{12b}, at each occurrence, is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-2 R^{12e}, and a $(CH_2)_r$-5-6 membered heterocyclic

system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{12e};

R^{12d}, at each occurrence, is independently selected from H, methyl, $-CF_3$, C_{2-4} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, a C_{3-6} carbocyclic residue substituted with 0-3 R^{12e}, and a $(CH_2)_r$-5-6 membered heterocyclic system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^{12e};

R^{12e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-6} cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, $-O-C_{1-6}$ alkyl, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{12f}R^{12f}$, and $(CH_2)_r$phenyl;

R^{12f}, at each occurrence, is selected from H, C_{1-6} alkyl, and C_{3-6} cycloalkyl;

~~R^{13}, at each occurrence, is independently selected from methyl, C_{2-4} alkyl substituted with 0-1 R^{13b};~~

~~R^{13b} is selected from $-OH$, $-SH$, $-NR^{13e}R^{13e}$, $-C(O)NR^{13e}R^{13e}$, and $-NHC(O)R^{13e}$;~~

~~R^{13e} is selected from H, C_{1-4} alkyl and C_{3-6} cycloalkyl;~~

n is selected from 1 and 2;

AMENDMENTS TO THE CLAIMS

~~m is selected from 0 and 1;~~

p, at each occurrence, is independently selected from 0, 1, and 2;

q, at each occurrence, is independently selected from 1, 2, 3, and 4;

r, at each occurrence, is independently selected from 0, 1, 2, 3, and 4;

~~s, at each occurrence, is independently selected from 0 and 1; and~~

t, at each occurrence, is independently selected from 2, 3, and 4.

3. (AMENDED) The compound of claim 2, wherein:

R^{10} and R^{10a} are H; and

~~m is 0;~~

n is 1~~; and~~

~~s is 0~~.

4. (AMENDED) The compound of claim 3, wherein:

AMENDMENTS TO THE CLAIMS

ring B is ~~selected from~~



~~and~~ ring B being optionally substituted with 0-1 R^5; and

R^{11} and R^{12} are H.

5. (ORIGINAL) The compound of claim 4, wherein:

R^5, at each occurrence, is independently selected from H, C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CRR)_rOH$, $(CRR)_rSH$, $(CRR)_rOR^{5d}$, $(CRR)_rSR^{5d}$, $(CRR)_rNR^{5a}R^{5a}$, $(CRR)_rC(O)OH$, $(CRR)_rC(O)R^{5b}$, $(CRR)_rC(O)NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}C(O)R^{5b}$, $(CRR)_rNR^{5a}C(O)OR^{5d}$, $(CRR)_rOC(O)NR^{5a}R^{5a}$, $(CHR)_rNR^{5a}C(O)NR^{5a}R^{5a}$, $CRR(CRR)_rNR^{5a}C(O)H$, $(CRR)_rC(O)OR^{5b}$, $(CRR)_rOC(O)R^{5b}$, $(CRR)_rS(O)_pR^{5b}$, $(CRR)_rS(O)_2NR^{5a}R^{5a}$, $(CRR)_rNR^{5a}S(O)_2R^{5b}$, and C_{1-6} haloalkyl;

R^{5a}, at each occurrence, is independently selected from H, methyl, C_{1-6} alkyl substituted with 0-2 R^{5e} wherein the alkyl is selected from ethyl, propyl,

i-propyl, butyl, i-butyl, pentyl, hexyl, C_3 alkenyl substituted with 0-1 R^{5e}, wherein the alkenyl is selected from allyl, C_3 alkynyl substituted with 0-1 R^{5e} wherein the alkynyl is selected from propynyl, and a $(CH_2)_r$-C_{3-4} carbocyclic residue substituted with 0-5 R^{5e}, wherein the carbocyclic residue is selected from cyclopropyl, and cyclobutyl;

R^{5b}, at each occurrence, is selected from C_{1-6} alkyl substituted with 0-2 R^{5e}, wherein the alkyl is selected from methyl, ethyl, propyl, i-propyl, butyl, i-butyl, pentyl, and hexyl, a $(CH_2)_r$-C_{3-4} carbocyclic residue substituted with 0-2 R^{5e}, wherein the carbocyclic residue is selected from cyclopropyl, and cyclobutyl; and

R^{5d}, at each occurrence, is selected from methyl, CF_3, C_{2-6} alkyl substituted with 0-2 R^{5e}, wherein the alkyl is selected from methyl, ethyl, propyl, i-propyl, butyl, i-butyl, pentyl, and hexyl, C_{3-8} alkenyl, C_{3-8} alkynyl, and a C_{3-10} carbocyclic residue substituted with 0-3 R^{5e}.

6. (ORIGINAL) The compound of claim 5, wherein:

R^4 is selected from H, C_{1-6} alkyl, C_{3-8} alkenyl, C_{3-8} alkynyl, $(CRR)_qOH$, $(CRR)_tSH$, $(CRR)_tOR^{4d}$, $(CRR)_tSR^{4d}$, $(CRR)_tNR^{4a}R^{4a}$, $(CRR)_qC(O)OH$, $(CRR)_rC(O)R^{4b}$, $(CRR)_rC(O)NR^{4a}R^{4a}$, $(CRR)_tNR^{4a}C(O)R^{4b}$,

AMENDMENTS TO THE CLAIMS

$(CRR)_tOC(O)NR^{4a}R^{4a}$, $(CRR)_tNR^{4a}C(O)OR^{4d}$, $(CRR)_tNR^{4a}C(O)R^{4b}$, $(CRR)_rC(O)OR^{4b}$, $(CRR)_tOC(O)R^{4b}$, $(CRR)_rS(O)_pR^{4b}$, $(CRR)_rS(O)_2NR^{4a}R^{4a}$, $(CRR)_rNR^{4a}S(O)_2R^{4b}$;

R, at each occurrence, is independently selected from H, methyl, ethyl, propyl, allyl, propynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, and $(CH_2)_r$phenyl substituted with R^{6e};

R^5, at each occurrence, is independently selected from H, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, allyl, propynyl, $(CH_2)_rOH$, $(CH_2)_rOR^{5d}$, $(CH_2)_rNR^{5a}R^{5a}$, $(CH_2)_rC(O)OH$, $(CH_2)_rC(O)R^{5b}$, $(CH_2)_rC(O)NR^{5a}R^{5a}$, $(CH_2)_rNR^{5a}C(O)R^{5b}$, $(CH_2)_rOC(O)NR^{5a}R^{5a}$, $(CH_2)_rNR^{5a}C(O)OR^{5d}$, $(CH_2)_rNR^{5a}C(O)R^{5b}$, $(CH_2)_rC(O)OR^{5b}$, $(CH_2)_rOC(O)R^{5b}$, $(CH_2)_rNR^{5a}S(O)_2R^{5b}$, and C_{1-6} haloalkyl;

R^{5a}, at each occurrence, is independently selected from H, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, pentyl, hexyl, cyclopropyl, and cyclobutyl; and

r, at each occurrence, is selected from 0, 1, and 2.

7. (ORIGINAL) The compound of claim 6, wherein:

R^1 is selected from phenyl substituted with 0-2 R^6, naphthyl substituted with 0-2R^6, and a 5-10

membered heteroaryl system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^6 wherein the heteroaryl is selected from indolyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalonyl, cinnolinyl, furanyl, imidazolyl, indazolyl, indolyl, isoquinolinyl isothiazolyl, isoxazolyl, oxazolyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridyl, pyridinyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolinyl, thiazolyl, thienyl, and tetrazolyl;

R^2 is selected from phenyl substituted with 0-2 R^7, and a 5-10 membered heteroaryl system containing 1-4 heteroatoms selected from N, O, and S, substituted with 0-3 R^7 wherein the heteroaryl is selected from indolyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalonyl, cinnolinyl, furanyl, imidazolyl, indazolyl, indolyl, isoquinolinyl isothiazolyl, isoxazolyl, oxazolyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridyl, pyridinyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolinyl, thiazolyl, thienyl, and tetrazolyl;

R^4 is selected from H, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, allyl, propynyl, $(CRR)_qOH$, $(CRR)_tSH$, $(CRR)_tOR^{4d}$, $(CRR)_tSR^{4d}$, $(CRR)_tNR^{4a}R^{4a}$, $(CRR)_qC(O)OH$, $(CRR)_rC(O)R^{4b}$, $(CRR)_rC(O)NR^{4a}R^{4a}$, $(CRR)_tNR^{4a}C(O)R^{4b}$, $(CRR)_tOC(O)NR^{4a}R^{4a}$,

$(CRR)_tNR^{4a}C(O)OR^{4d}$, $(CRR)_tNR^{4a}C(O)R^{4b}$, $(CRR)_rC(O)OR^{4b}$, $(CRR)_tOC(O)R^{4b}$, $(CRR)_rS(O)_pR^{4b}$, $(CRR)_rS(O)_2NR^{4a}R^{4a}$, $(CRR)_rNR^{4a}S(O)_2R^{4b}$;

R^{4a}, at each occurrence, is independently selected from H, methyl substituted with 0-1 R^{4c}, C_{2-6} alkyl substituted with 0-3 R^{4e} wherein C_{2-6} is selected from ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl and hexyl, and a $(CH_2)_r$-C_{3-6} carbocyclic residue substituted with 0-4 R^{4e} wherein the carbocyclic residue is selected from cyclopropyl, cyclohexyl, and phenyl;

R^{4b} is selected from H, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, and cyclopropyl;

R^{4d} is selected from methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, and cyclopropyl;

R^8 is selected from H, methyl, ethyl, propyl, i-propyl, and cyclopropyl; and

R^9 is selected from H, methyl, ethyl, propyl, i-propyl, and cyclopropyl, and CH_2-R^1.

8. (ORIGINAL) The compound of claim 7, wherein:

R^6, at each occurrence, is selected from C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CRR)_rC_{3-6}$ cycloalkyl, Cl, Br, I, F, NO_2, CN, $(CRR)_rNR^{6a}R^{6a}$, $(CRR)_rOH$,

$(CRR)_rO(CRR)_rR^{6d}$, $(CRR)_rSH$, $(CRR)_rC(O)H$, $(CRR)_rS(CRR)_rR^{6d}$, $(CRR)_rC(O)OH$, $(CRR)_rC(O)(CRR)_rR^{6b}$, $(CRR)_rC(O)NR^{6a}R^{6a}$, $(CRR)_rNR^{6f}C(O)(CRR)_rR^{6b}$, $(CRR)_rC(O)O(CRR)_rR^{6d}$, $(CRR)_rNR^{6a}C(O)NR^{6a}R^{6a}$, $(CRR)_rNR^{6a}C(S)NR^{6a}R^{6a}$, $(CRR)_rOC(O)(CRR)_rR^{6b}$, $(CRR)_rS(O)_p(CRR)_rR^{6b}$, $(CRR)_rS(O)_2NR^{6a}R^{6a}$, $(CRR)_rNR^{6f}S(O)_2(CRR)_rR^{6b}$, $(CRR)_rNR^{6f}S(O)_2$ $NR^{6a}R^{6a}$, C_{1-6} haloalkyl, and $(CRR)_r$phenyl substituted with 0-3 R^{6e};

R^{6a}, at each occurrence, is independently selected from H, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, hexyl, cyclopropyl and phenyl;

R^{6b}, at each occurrence, is selected from methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, hexyl, cyclopropyl, and phenyl;

R^{6d}, at each occurrence, is selected from methyl, CF_3, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, hexyl, cyclopropyl, and phenyl;

R^{6e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{6f}R^{6f}$, and $(CH_2)_r$phenyl;

R^{6f}, at each occurrence, is selected from H, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, hexyl, cyclopropyl, and phenyl;

R^7 is selected from methyl, ethyl, propyl, i-propyl, butyl, i-butyl, s- butyl, t-butyl, pentyl, hexyl, $(CRR)_rC_{3-6}$ cycloalkyl, Cl, Br, I, F, NO_2, CN, $(CRR)_rNR^{7a}R^{7a}$, $(CRR)_rOH$, $(CRR)_rO(CH)_rR^{7d}$, $(CRR)_rSH$, $(CRR)_rC(O)H$, $(CRR)_rS(CRR)_rR^{7d}$, $(CRR)_rC(O)OH$, $(CRR)_rC(O)(CRR)_rR^{7b}$, $(CRR)_rC(O)NR^{7a}R^{7a}$, $(CRR)_rNR^{7f}C(O)(CRR)_rR^{7b}$, $(CRR)_rC(O)O(CRR)_rR^{7d}$, $(CRR)_rOC(O)(CRR)_rR^{7b}$, $(CRR)_rNR^{7a}C(O)NR^{7a}R^{7a}$, $(CRR)_rNR^{7a}C(O)O(CRR)_rR^{7d}$, $(CRR)_rS(O)_p(CRR)_rR^{7b}$, $(CRR)_rS(O)_2NR^{7a}R^{7a}$, $(CRR)_rNR^{7f}S(O)_2(CRR)_rR^{7b}$, C_{1-6} haloalkyl, and $(CRR)_r$phenyl substituted with 0-3 R^{7e};

R^{7a}, at each occurrence, is selected from H, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, hexyl,, prop-2-enyl, 2-methyl-2-propenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, CH_2cyclopropyl, and benzyl;

R^{7b}, at each occurrence, is selected from methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, hexyl, cyclopropyl, cyclopentyl, CH_2-cyclopentyl, cyclohexyl, CH_2-cyclohexyl, CF_3, pyrrolidinyl, morpholinyl, and azetidinyl;

R^{7d}, at each occurrence, is selected from methyl, CF_3, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, hexyl, and cyclopropyl;

R^{7e}, at each occurrence, is selected from C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, $(CH_2)_rC_{3-6}$ cycloalkyl, Cl, F, Br, I, CN, NO_2, $(CF_2)_rCF_3$, $(CH_2)_rOC_{1-5}$ alkyl, OH, SH, $(CH_2)_rSC_{1-5}$ alkyl, $(CH_2)_rNR^{7f}R^{7f}$, and $(CH_2)_r$phenyl;

R^{7f}, at each occurrence, is selected from H, methyl, ethyl, propyl, i-propyl, butyl, i-butyl, t-butyl, pentyl, hexyl, cyclopropyl, and phenyl; and

r is 0 or 1.

9. (ORIGINAL) The compound of claim 8, wherein

R^7 is selected from methyl, ethyl, propyl, i-propyl, butyl, i-butyl, s-butyl, pentyl, hexyl, Cl, Br, I, F, NO_2, $NR^{7a}R^{7a}$, $NHC(O)NHR^{7a}$, $NR^{7a}C(O)R^{7b}$, $NR^{7a}C(O)OR^{7d}$, CF_3, OCF_3, $C(O)R^{7b}$, $NR^{7f}C(O)NR^{7a}R^{7a}$, $NHS(O)_2R^{7b}$,



,



, and



.

AMENDMENTS TO THE CLAIMS

10. (AMENDED) The compound of claim 9, wherein

~~ring B is selected from~~  ~~and~~

Z is -C(O)-;

~~R^{1a} and R^{1b} are selected from H and methyl, or alternatively, R^{1a} and R^{1b} are taken together to form =O;~~

R^1 is selected from a C_{6-10} aryl group substituted with 0-3 R^6 wherein the aryl group is selected from phenyl and naphthyl, and a 5-10 membered heteroaryl system containing 1-4 heteroatoms selected from N and O, substituted with 0-3 R^6 wherein the heteroaryl system is selected from furyl, indolyl, and benzotriazolyl;

R^2 is phenyl substituted with 0-1 R^7;

R^4 is selected from H, methyl, ethyl, propyl, i-propyl, butyl, I-butyl, t-butyl, pentyl, hexyl, and $(CH_2)_r C(O)R^{4b}$;

R^6 is selected from methyl, ethyl, propyl, i-propyl, butyl, F, Cl, Br, I, NO_2, CN, $O(CH_2)_rR^{6d}$, C(O)H, SR^{6d}, $NR^{6a}R^{6a}$, $OC(O)R^{6b}$, $S(O)_pR^{6b}$, $(CHR')_rS(O)_2NR^{6a}R^{6a}$, CF_3;

R^{6a} is H methyl, or ethyl;

R^{6b} is H or methyl;

R^{6d} is methyl, phenyl, CF_3, and (CH_2)-phenyl;

R^9 is selected from H, methyl, and (CH_2)-R^1; and

r is 0 or 1.

11. (AMENDED) The compound of claim 1, wherein the compound is selected from:

N-[2-[[(1S,2S)-2-[[(4-Chlorophenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[[(2,4-Dimethylphenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[[(2,4,6-Trimethylphenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

AMENDMENTS TO THE CLAIMS

N-[2-[[(1S,2S)-2-[[(4-Benzyloxyphenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[[(2,4-Difluorophenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[[(2-Chloro-4-fluorophenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[[(2-Trifluoromethyl-4-fluorophenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[[(2,4-Dichlorophenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[[(2-Fluoro-6-trifluoromethylphenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[[(2-Chloro-5-trifluoromethylphenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[[(1-Naphthyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[bis(3-furylmethyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[(2,4-Dimethylbenzyl)(methyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(1S,2S)-2-[(4-Chlorobenzyl)(methyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(cis)-2-[[(2,4-Dimethylphenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(cis)-2-[[(4-Chlorophenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(cis)-2-[[(4-Nitrophenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(cis)-2-[[(4-Isopropylphenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(cis)-2-[[(4-Trifluorophenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

AMENDMENTS TO THE CLAIMS

N-[2-[[(cis)-2-[[(4-Trifluoromethoxyphenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(cis)-2-[[(4-Phenoxyphenyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(cis)-2-[[(1-Naphthyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(cis)-2-[[(2-Naphthyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

N-[2-[[(cis)-2-[[(3-Indolyl)methyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

~~N-[2-[[(cis)-2-[[1-(4-Chlorophenyl)ethyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

N-[2-[[(cis)-2-[Bis(3-furylmethyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;

~~N-[2-[[(1S,2R)-2-[(4-Chlorobenzoyl)amino]cyclopentyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

AMENDMENTS TO THE CLAIMS

~~N-[2-[[(1S,2R)-2-[(4-(Methylthio)benzoyl)amino]cyclopentyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(1S,2R)-2-[(4-(Methylsulfonyl)benzoyl)amino]cyclopentyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(1S,2R)-2-[(4-Iodobenzoyl)amino]cyclopentyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(1S,2R)-2-[(4-(Aminosulfonyl)benzoyl)amino]cyclopentyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(1S,2R)-2-[[(4-Chlorophenyl)methyl]amino]cyclopentyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(1S,2R)-2-[[(2,4-Dimethylphenyl)methyl]amino]cyclopentyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(1S,2R)-2-[[(4-Methylphenyl)methyl]amino]cyclopentyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Chlorobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

AMENDMENTS TO THE CLAIMS

~~N-[2-[[(cis)-2-[(4-Methylbenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Fluorobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[Benzoylamino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Bromobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Phenoxybenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Trifluoromethylbenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(5-Benzotriazolecarbonyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Iodobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

AMENDMENTS TO THE CLAIMS

~~N-[2-[[(cis)-2-[(4-Cyanobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Trifluoromethoxybenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Formylbenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Carbomethoxybenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Nitrobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Aminobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Methoxybenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Methylthiobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Methylsulfonylbenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Aminosulfonylbenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Isopropylbenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Phenylthiobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-(N,N-diethylsulfamoyl)benzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[(4-Trifluoromethylthiobenzoyl)amino]cyclohexyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[[(4-Chlorophenyl)methyl]amino]cyclopropyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[[(3,4-Dimethylphenyl)methyl]amino]cyclopropyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~N-[2-[[(cis)-2-[[(4-Methylphenyl)methyl]amino]cyclopropyl]amino]-2-oxoethyl]-3-(trifluoromethyl)benzamide;~~

~~2-Amino-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-iodobenzamide;~~

~~2-Amino-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-chlorobenzamide;~~

~~N-[2-[[(cis)-2-[[4-(Aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-chlorobenzamide;~~

~~N-[2-[[(cis)-2-[[4-(Aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-3-trifluoromethoxybenzamide;~~

~~Tert-butyl 2-[([2-[((cis)-2-[[4-(aminosulfonyl)benzoyl]amino}cyclohexyl)amino]-2-oxoethyl}amino)carbonyl]-4-(trifluoromethyl)phenylcarbamate;~~

~~2-Amino-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethylbenzamide trifluoroacetate;~~

AMENDMENTS TO THE CLAIMS

~~4-(Aminosulfonyl)-N-((cis)-2-{[({[2-(trifluoromethyl)anilino]carbonyl}amino)acetyl]amino}cyclohexyl)benzamide;~~

~~4-(Aminosulfonyl)-N-{(cis)-2-[({[(3-chlorophenyl)sulfonyl]amino}acetyl)amino]cyclohexyl}benzamide;~~

~~Ethyl 2-[({2-[((cis)-2-{[4-(aminosulfonyl)benzoyl]amino}cyclohexyl)amino]-2-oxoethyl}amino)carbonyl]-4-(iodo)phenylcarbamate;~~

AMENDMENTS TO THE CLAIMS

~~Methyl 2-[({2-[((cis)-2-{[4-(aminosulfonyl)benzoyl]amino}cyclohexyl)amino]-2-oxoethyl}amino)carbonyl]-4-(iodo)phenylcarbamate;~~

~~Tert-butyl N-Methyl-2-[({2-[((cis)-2-{[4-(aminosulfonyl)benzoyl]amino}cyclohexyl)amino]-2-oxoethyl}amino)carbonyl]-4-(trifluoromethyl)phenylcarbamate;~~

~~Ethyl 2-[({2-[((cis)-2-{[4-(aminosulfonyl)benzoyl]amino}cyclohexyl)amino]-2-oxoethyl}amino)carbonyl]-4-(trifluoromethyl)phenylcarbamate;~~

~~2-(Benzylamino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-(Ethylamino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-(Methylamino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-Amino-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-bromo benzamide;~~

~~Tert-butyl 2-[({2-[((cis)-2-{[4-(aminosulfonyl)benzoyl]amino}cyclohexyl)amino]-2-~~

~~oxoethyl]amino)carbonyl]-4-(trifluoromethoxy)phenylcarbamate;~~

~~2-Amino-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethoxy benzamide;~~

~~2-(Allylamino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((2-methyl-2-propenyl)amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-(cyclopropylmethylene)amino-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-(butyl)amino-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-(propyl)amino-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-(propyl)amino-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

AMENDMENTS TO THE CLAIMS

~~2-((2-methyl-2-propyl)amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl-benzamide;~~

~~2-((aminocarbonyl)amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl-benzamide;~~

~~2-(acetylamino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl-benzamide;~~

~~2-(Methylamino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-iodomethyl-benzamide;~~

~~2-(Ethylamino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-iodomethyl-benzamide;~~

~~2-(Trifluoroacetylamino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-iodomethyl-benzamide;~~

~~2-(amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-nitro-benzamide;~~

~~Iso-propyl-2-[({2-[((cis)-2-{[4-(aminosulfonyl)benzoyl]amino}cyclohexyl)amino]-2-oxoethyl}amino)carbonyl]-4-(iodo)phenylcarbamate;~~

AMENDMENTS TO THE CLAIMS

~~Tert-butyl 2-[({2-[((cis)-2-{[4-(aminosulfonyl)benzoyl]amino}cyclohexyl)amino]-2-oxoethyl}amino)carbonyl]-4-(iodo)phenylcarbamate;~~

~~2-(amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-3,5-dinitro benzamide;~~

~~2-((Isopropylaminocarbonyl)amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((cyclohexylcarbonyl)amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Cyclopentylmethylenecarbonyl)amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((cyclohexylcarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylsulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((cyclohexylcarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Isopropylaminocarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

AMENDMENTS TO THE CLAIMS

~~2-((Isopropylaminocarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylsulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Methylsulfonyl)amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Aminocarbonyl)amino)-N-[2-[[(cis)-2-[[4-(aminosulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Allyl)amino)-N-[2-[[(cis)-2-[[4-(methylsulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Allyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((2-Methyl-2-propenyl)amino)-N-[2-[[(cis)-2-[[4-(methylsulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((2-methyl-2-propenyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Propyl)amino)-N-[2-[[(cis)-2-[[4-(methylsulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

AMENDMENTS TO THE CLAIMS

~~2-((Propyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((2-Methylpropyl)amino)-N-[2-[[(cis)-2-[[4-(methylsulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((2-Methylpropyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Butyl)amino)-N-[2-[[(cis)-2-[[4-(methylsulfonyl)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Butyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Ethylaminocarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Allylaminocarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

AMENDMENTS TO THE CLAIMS

~~2-((Iso-butylaminocarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Cyclopentylaminocarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Tert-butoxycarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Iso-propoxycarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Ethoxycarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Pyrrolidinylcarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Morpholinylcarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

~~2-((Azetidinylcarbonyl)amino)-N-[2-[[(cis)-2-[[4-(methylthio)benzoyl]amino]cyclohexyl]amino]-2-oxoethyl]-5-trifluoromethyl benzamide;~~

AMENDMENTS TO THE CLAIMS

~~2-{[1-Pyrrolidinylcarbonyl]amino}-*N*-{2-[((cis)-4-{[4-(methylthio)benzyl]amino}tetrahydro-2*H*-pyran-3-yl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~2-{[1-Azetidinylcarbonyl]amino}-*N*-{2-[((cis)-4-{[4-(methylthio)benzyl]amino}tetrahydro-2*H*-pyran-3-yl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~2-{[1-Azetidinylcarbonyl]amino}-*N*-{2-[((cis)-4-{[4-(methoxy)benzyl]amino}tetrahydro-2*H*-pyran-3-yl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~1-(4-Methylthiobenzoylamino)-2-[2-(2-amino-5-trifluoromethylbenzoylamino)-acetylamino]-4-aminocyclohexane~~;

~~[2-({[5-benzyloxycarbonylamino-2-(4-methylthio-benzoylamino)cyclohexylcarbamoyl]-methyl}carbamoyl)-4-trifluoromethylphenyl]-carbamic acid tert-butyl ester~~;

~~{4-(4-Methylthiobenzoylamino)-3-[2-(3-trifluoromethylbenzoylamino)-acetylamino]-4-aminocyclohexane;~~

~~{4-(4-methylthiobenzoylamino)-3-[2-(3-trifluoromethylbenzoylamino)acetylamino]-cyclohexyl}carbamic acid benzyl ester;~~

AMENDMENTS TO THE CLAIMS

~~1-(4-Methanesulfonylbenzoylamino)-2-[2-(3-trifluoromethylbenzoylamino)-acetylamino]cyclohexyl-4-aminocyclohexane;~~

~~1-(4-Methylthiobenzoylamino)-2-[2-(2-amino-5-trifluoromethylbenzoylamino)acetylamino]-4-(2-propylamino)cyclohexane;~~

~~1-(4-Methylthiobenzoylamino)-2-[2-(2-amino-5-trifluoromethylbenzoylamino)acetylamino]-4-(3-methylureido)cyclohexane;~~

~~1-(4-Methylthiobenzoylamino)-2-[2-(3-trifluoromethylbenzoylamino)acetylamino]6-aminocyclohexane;~~

~~1-(4-Methylthiobenzoylamino)-2-[2-(3-trifluoromethylbenzoylamino)acetylamino]6-(2-propylamino)cyclohexane;~~

~~1-(4-Methylthio-benzoylamino)-2-[2-(2-Amino-5-trifluoromethyl-benzoylamino)-acetylamino]-4-aminocyclohexane;~~

~~4-(4-Methylthiobenzoylamino)-3-[2-(3-trifluoromethylbenzoylamino)acetylamino]-4-(2-propylamino)-cyclohexane;~~

~~1-(4-Methylthiobenzoylamino)-2-[2-(3-trifluoromethylbenzoylamino)acetylamino]-5-aminocyclohexane;~~

2-Amino-N-({2-[(4-methylthiophenylamino)methyl]cyclohexylcarbamoyl}-methyl)-5-(trifluoromethyl)benzamide;

2-Isopropylamino-N-{[(cis)2-(4-methylthiobenzylamino)-cyclohexylcarbamoyl]-methyl}-5-trifluoromethyl-benzamide;

2-(3-Isopropylureido)-N-{[2-(4-methylthiobenzylamino)cyclohexylcarbamoyl]-methyl}-5-trifluoromethylbenzamide;

2-(3-Morpholinylureido)-N-{[2-(4-methylthiobenzylamino)cyclohexylcarbamoyl]-methyl}-5-trifluoromethylbenzamide;

~~2-Amino-N-({2-(cis)-[3-(4-methylthiophenyl)ureido]cyclohexylcarbamoyl}methyl)-5-trifluoromethyl-benzamide;~~

~~{2-[({2-(Cis)-[3-(4-methanesulfonylphenyl)ureido]cyclohexylcarbamoyl}methyl)-carbamoyl]-4-trifluoromethylphenyl}-carbamic acid tert-butyl ester;~~

~~2-amino-*N*-{2-[((3*S*,4*R*)-4-{[4-(methylthio)benzyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~2-Amino-*N*-{2-[((3*R*,4*S*)-4-{[4-(methylthio)benzyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

AMENDMENTS TO THE CLAIMS

~~2-amino-*N*-{2-[((cis)-4-{[4-(methylthio)benzoyl]amino}-1-methyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~*N*-{2-[((cis)-4-{[4-chlorobenzyl]amino}-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

~~*N*-{2-[((cis)-4-{[4-(methylthio)benzyl]amino}-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

~~2-Amino-*N*-{2-[((cis)-4-{[4-chlorobenzyl]amino}-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~2-Amino-*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~2-Amino-*N*-{2-[((cis)-4-{[4-ethylthiobenzyl]amino}-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-1-methyl-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

AMENDMENTS TO THE CLAIMS

~~*N*-{2-[((cis)-4-{bis[4-methylthiobenzyl]amino}-1-methyl-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

~~2-Amino-*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-1-methyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-1-acetyl-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

~~2-Amino-*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-1-butyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~2-Cyclohexylamino-*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~2-Iso-propylamino-*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~2-(Pyrrolidinylcarbonyl)amino-*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

AMENDMENTS TO THE CLAIMS

~~2-(Methylaminocarbonyl)amino-*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~3-Amino-*N*-{2-[((cis)-4-{[4-methylthiobenzyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~*N*-{2-[((cis)-4-{[4-aminosulfonylbenzoyl]amino}-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

~~*N*-{2-[((cis)-4-{[4-methylsulfonylbenzoyl]amino}-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

~~2-Amino-*N*-{2-[((cis)-4-{[4-(methylthio)benzoyl]amino}-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~*N*-{2-[((cis)-4-{[4-methylthiobenzoyl]amino}-1-methyl-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

~~*N*-{2-[((cis)-4-{[4-methylthiobenzoyl]amino}-1-acetyl-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

AMENDMENTS TO THE CLAIMS

~~2-Amino-*N*-{2-[((cis)-4-{[4-methylthiobenzoyl]amino}-1-butyl-3-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

~~2-Cyclohexylamino-*N*-{2-[((cis)-4-{[4-methylthiobenzoyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~2-Iso-propylamino-*N*-{2-[((cis)-4-{[4-methylthiobenzoyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~3-Amino-*N*-{2-[((cis)-4-{[4-methylthiobenzoyl]amino}-1-propyl-3-piperidinyl)amino]-2-oxoethyl}-5-(trifluoromethyl)benzamide;~~

~~*N*-{2-[((cis)-3-{[4-(aminosulfonyl)benzoyl]amino}-4-piperidinyl)amino]-2-oxoethyl}-3-(trifluoromethyl)benzamide;~~

N-{[4-Dimethylamino-2-(4-methylsulfanyl-benzylamino)-cyclohexylcarbamoyl]-methyl}-3-trifluoromethyl-benzamide trifluoroacetate;

N-{[2-(4-Chloro-benzylamino)-4-dimethylamino-cyclohexylcarbamoyl]-methyl}-3-trifluoromethyl-benzamide trifluoroacetate;

N-{[4-Dimethylamino-2-(4-methoxy-benzylamino)-cyclohexylcarbamoyl]-methyl}-3-trifluoromethyl-benzamide trifluoroacetate; and

N-{[4-Dimethylamino-2-(4-methyl-benzylamino)-cyclohexylcarbamoyl]-methyl}-3-trifluoromethyl-benzamide trifluoroacetate.

12. (ORIGINAL) A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claim 1.

13. (ORIGINAL) A method for modulation of chemokine or chemokine receptor activity comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1.

14. (ORIGINAL) A method for modulation of MCP-1, MCP-2, MCP-3 and MCP-4, and MCP-5 activity that is mediated by the CCR2 receptor comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1.

15. (ORIGINAL) A method for modulation of MCP-1 activity comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1.

16. (AMENDED) A method for treating ~~or preventing~~ disorders, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1, said disorders being selected from osteoarthritis, aneurism, fever, cardiovascular effects, Crohn's disease, congestive heart failure, autoimmune diseases, HIV-infection, HIV-associated dementia, psoriasis, idiopathic pulmonary fibrosis, transplant arteriosclerosis, physically- or chemically-induced brain trauma, inflammatory bowel disease, alveolitis, colitis, systemic lupus erythematosus, nephrotoxic serum nephritis, glomerularnephritis, asthma, multiple sclerosis, artherosclerosis, and rheumatoid arthritis.

17. (AMENDED) The method for treating ~~or preventing~~ disorders, of claim 16, wherein said disorders being selected from psoriasis, idiopathic pulmonary fibrosis, transplant arteriosclerosis, physically- or chemically-induced brain trauma, inflammatory bowel disease, alveolitis, colitis, systemic lupus erythematosus, nephrotoxic serum nephritis, glomerularnephritis, asthma, multiple sclerosis, artherosclerosis, and rheumatoid arthritis.

18. (AMENDED) The method for treating ~~or preventing~~ disorders, of claim 17, wherein said disorders being selected from alveolitis, colitis, systemic lupus erythematosus, nephrotoxic serum nephritis, glomerularnephritis, asthma, multiple sclerosis, artherosclerosis, and rheumatoid arthritis.

19. (AMENDED) The method for treating ~~or preventing~~ disorders, of claim 18, wherein said disorders being selected from asthma, multiple sclerosis, artherosclerosis, and rheumatoid arthritis.

20. (AMENDED) A method for treating ~~or preventing~~ rheumatoid arthritis, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1.

21. (AMENDED) A method for treating ~~or preventing~~ multiple sclerosis, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1.

22. (AMENDED) A method for treating ~~or preventing~~ atherosclerosis, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1.

23. (AMENDED) A method for treating ~~or preventing~~ asthma, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1.

24. (AMENDED) A method for treating ~~or preventing~~ inflammatory diseases, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1.

25. (ORIGINAL) A method for modulation of CCR2 activity comprising administering to a patient in need

thereof a therapeutically effective amount of a compound of claim 1.

26. (NEW) A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claim 8.

27. (NEW) A method for modulation of MCP-1 activity comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 8.

28. (NEW) The method for treating ~~or preventing~~ disorders, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 8, wherein said disorders being selected from asthma, multiple sclerosis, artherosclerosis, and rheumatoid arthritis.

29. (NEW) A method for treating ~~or preventing~~ rheumatoid arthritis, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 8.